

## CONTENTS TO VOLUME 128

## Hypothesis

- Peroxisomal respiration and energy conservation: Possible relationships between heat production, thermosmosis and conformational changes, *D. J. Hryb* 1

## Research Letters

- Effect of the chemotactic peptide on the subsequent superoxide-releasing response in human polymorphonuclear leukocytes, *S. Kitagawa, F. Takaku* 5
- Heparin-bound lipoprotein lipase is catalytically active and can be stimulated by apolipoprotein CII, *G. Bengtsson, T. Olivecrona* 9
- Further studies on subunit III of bovine procarboxypeptidase A. Structure and reactivity of the weakly functional active site, *C. Wicker, A. Puigserver* 13
- Labelling of the hydrophobic domain of the Na<sup>+</sup>,K<sup>+</sup>-ATPase, *C. Montecucco, R. Bisson, C. Gache, A. Johannsson* 17
- Immobilized D-glyceraldehyde-3-phosphate dehydrogenase can exist as a trimer, *L. I. Ashmarina, V. I. Mironetz, N. K. Nagradova* 22
- A novel location for two enzymes of de novo pyrimidine biosynthesis in trypanosomes and *Leishmania*, *D. J. Hammond, W. E. Gutteridge, F. R. Oppendoerfer* 27
- 3-Methyl-5-(4'-oxo-2',6'-trimethylcyclohex-2'-en-1'-yl)-2,4 pentadienoic acid, a putative precursor of abscisic acid from *Cercospora roscicola*, *S. J. Neill, R. Horgan, T. S. Lee, D. C. Walton* 30
- Reversal of heparin inhibition of nuclear protein kinase NII by polyamines and histones, *T. Hara, K. Takahashi, H. Endo* 33
- Study of 'difficult peptides' from *Paracoccus* cytochrome c-550 and a dolphin cytochrome c. Fast atom bombardment; a new method for molecular mass and sequence determination of peptides, *D. H. Williams, G. Bojesen, A. D. Auffret, L. C. E. Taylor* 37
- Heterogeneity of nucleosomes in genetically inactive cells, *I. A. Zalenskaya, A. O. Zalensky, E. O. Zalenskaya, V. I. Vorob'ev* 40
- Study on the Met-tRNA-binding activity in Krebs-2 cells infected with encephalomyocarditis virus, *A. M. Dizhoor, T. Yu. Ugarova* 43
- Topography of interaction of *Escherichia coli* RNA polymerase subunits with *lac* UV5 promoter, *A. Chenchick, R. Beabealashvili, A. Mirzabekov* 46
- (D-Ala<sup>2</sup>, N-Val<sup>5</sup>), (D-Ala<sup>2</sup>, Met<sup>5</sup>) Enkephalins and enkephalinamides: Correlation between biological activity and NMR features, *J. E. Gairin, R. Haran, H. Mazarguil, Y. Audigier, J. Cros* 51
- Effects of low frequency electrical stimulation on enzyme and isozyme patterns of dystrophic mouse muscle, *H. Reichmann, D. Pette, G. Vrbová* 55
- Cholera toxin can recouple hormone receptors that are uncoupled from adenylyl cyclase, *P. A. Insel, F. J. Darfler, M. S. Kennedy* 59
- Regulation of the degradation of <sup>125</sup>I-labeled glutamine synthetase introduced into cultured hepatoma cells by erythrocyte ghost-mediated injection, *A. Freikopf-Cassel, R. G. Kulka* 63
- Isolation and characterization of a  $\gamma_1$ -melanotropin-like peptide from bovine neurointermediate pituitary, *P. Böhlen, F. Esch, T. Shibasaki, A. Baird, N. Ling, R. Guillemin* 67
- Potential use of glycosphingolipids of human meconium for blood group chemotyping of single individuals, *K.-A. Karlsson, G. Larson* 71

|   |     |
|---|-----|
| Stimulation of rat fat cell phosphodiesterase by adenosine, T.-S. Teo, S.-O. Ooi, E. H. A. Wong   | 75  |
| Metabolism of macromolecular methyl groups in <i>Escherichia coli</i> : Whole cell NMR spectroscopy, F. Hayashi, J. G. Thompson, P. F. Agris  | 79  |
| Peptidyl-puromycin synthesis on polysomes from starving bacteria, J. C. Cortay, A. J. Cozzone   | 83  |
| Comparative effect of S-adenosyl-homocysteine (SAH) and sinefungin on tRNA-base methylation in whole cells and in vitro, M. Vedel, M. Robert-Géro   | 87  |
| Nucleodisome — A new repeat unit of chromatin revealed in nuclei of pigeon erythrocytes by DNase I digestion, A. T. Khachatryan, V. A. Pospelov, S. B. Svetlikova, V. I. Vorob'ev   | 90  |
| Inhibition of the induction of heat shock proteins in <i>Drosophila melanogaster</i> cells infected with insect picornaviruses, N. F. Moore, J. S. K. Pullin, B. Reavy  | 93  |
| Orientation of $\beta$ -carotene and retinal in lipid bilayers, L. B.-Å. Johansson, G. Lindblom, A. Wieslander, G. Arvidsson  | 97  |
| Effects of modifications of leghemoglobin structure on its ability to oxidize indole-3-acetic acid, U. Perttälä   | 100 |
| A monoclonal antibody inhibiting rat liver 5'-nucleotidase, K. Siddle, E. M. Bailyes, J. P. Luzio   | 103 |
| Structural requirement for the biological activity of serum thymic factor, A. Imaizumi, J. Gyotoku, S. Terada, E. Kimoto  | 108 |
| Photo-affinity inactivation of gonadotropin releasing hormone receptors, E. Hazum   | 111 |
| Testosterone-induced, sulfonamide-resistant carbonic anhydrase isozyme of rat livers indistinguishable from skeletal muscle carbonic anhydrase III, N. D. Carter, D. Hewett-Emmett, S. Jeffret, R. E. Tashian   | 114 |
| Identification of a second binding region on rabbit skeletal troponin-T for $\alpha$ -tropomyosin, J. R. Pearlstone, L. B. Smillie  | 119 |
| Trypsin-induced activation of the thiol esters in $\alpha_2$ -macroglobulin generates a short-lived intermediate ('nascent' $\alpha_2$ M) that can react rapidly to incorporate not only methylamine or putrescine but also proteins lacking proteinase activity, L. Sottrup-Jensen, T. E. Petersen, S. Magnusson           | 123 |
| Mechanism of proteinase complex formation with $\alpha_2$ -macroglobulin. Three modes of trypsin binding, L. Sottrup-Jensen, T. E. Petersen, S. Magnusson   | 127 |
| ADP-ribosylation of sarcolemma membrane proteins in the presence of cholera toxin and its influence on insulin-stimulated membrane protein kinase activity, O. Walaas, R. S. Horn, E. Lystad, A. Adler  | 133 |
| Insulin-dependent protein phosphorylation in membranes. Isolation and characterization of a phosphorylated proteolipid from sarcolemma, O. Walaas, K. Sletten, R. S. Horn, E. Lystad, A. Adler, A. R. Alertsen  | 137 |
| Purification of a phosphate carrier in pig heart mitochondria by affinity chromatography on mersalyl-ultrogel, S. Touraille, Y. Briand, R. Druand   | 142 |
| 3R,S-3-Hydroxy-3-methyl-4-carboxybutyl-CoA; specific inhibitor of 3-hydroxy-3-methylglutaryl-CoA reductase, T.-G. Nguyen, H. Aigner, H. Eggerer   | 145 |
| Generation of large and active membrane layers from thylakoid vesicles. Properties and possible application, H. T. Witt, D. DiFiore   | 149 |
| Physical parameters of <i>Escherichia coli</i> translational initiation factor 3 binding to poly(A), E. Wickstrom   | 154 |
| Aminoacylation of ethionine to rat liver tRNA <sup>Met</sup> and its incorporation into protein, M. J. Wilson, D. L. Hatfield, L. A. Poirier  | 157 |
| A study of the interaction of <i>Escherichia coli</i> initiation factor IF-2 with formyl-methionyl-tRNA <sup>Met</sup> by partial digestion with cobra venom ribonuclease, H. U. Petersen, T. A. Kruse, H. Worm-Leonhard, G. E. Siboska, B. F. C. Clark, A. S. Boutorin, P. Remy, J. P. Ebel, J. Dondon, M. Grunberg-Manago | 161 |
| Book Reviews and Booklist No. 30  | 167 |
| Notes to Authors  | 176 |

## Review Letter

- Initiation of eukaryotic protein synthesis, *A. A. M. Thomas, R. Benne, H. O. Voorma* 177

## Research Letters

- Interaction of cAMP receptor protein with the *ompA* gene, a gene for a major outer membrane protein of *Escherichia coli*, *R. N. Movva, P. Green, K. Nakamura, M. Inouye* 186
- Interaction between latent papovavirus genomes and the tumor promoter TPA, *P. Krieg, E. Amtmann, G. Sauer* 191
- Acetylation of histone H<sub>4</sub> in chicken erythrocyte and cuttle-fish testis chromatin, *D. Wouters-Tyrou, A. Martin-Ponthieu, P. Sautière, G. Biserte* 195
- Enzymatic synthesis of the coenzyme A ester of *o*-succinylbenzoic acid, an intermediate in menaquinone (vitamin K<sub>2</sub>) biosynthesis, *L. Heide, E. Leistner* 201
- Photochemical action spectra of bacterial *a*- and *o*-type oxidases using a dye laser, *C. Edwards, S. Beer, A. Siviram, B. Chance* 205
- Glycolipid formation in *Volvox carteri* f. *nagariensis*: Effects of tunicamycin and showdomycin, *Th. Müller, E. Bause, L. Jaenicke* 208
- Resolution of *Streptococcus mutans* glucosyltransferases into two components essential to water-insoluble glucan synthesis, *K. Fukushima, R. Motoda, K. Takada, T. Ikeda* 213
- Porcine pancreatic procolipase and its trypsin activated form. Lipid binding and lipase activation on monomolecular films, *T. Wieloch, B. Borgström, G. Piéroni, F. Pattus, R. Verger* 217
- Age-associated accumulation of altered FDP aldolase B in mice. Conditions of detection and determination of aldolase half-life in young and old animals, *A. Z. Reznick, L. Lavie, H. E. Gershon, D. Gershon* 221
- Comparative patterns of 'in vitro' oxidative hemolysis of normal and glucose 6-phosphate dehydrogenase (G6PD)-deficient erythrocytes, *U. Benatti, A. Morelli, T. Meloni, B. Sparatore, F. Salamino, M. Michetti, E. Melloni, S. Pontremoli, A. de Flora* 225
- Effect of adrenalectomy and cortisol treatment on dihydroxyacetone phosphate acyltransferase activity in rat liver, *E. J. Bates, E. D. Saggerson* 230
- Differential distribution of tropomyosin subunits in fast and slow rat muscles and its changes in long-term denervated hemidiaphragm, *U. Carraro, C. Catani, L. Dalla Libera, M. Vascon, G. Xanella* 233
- CoA-dependent cleavage of arachidonic acid from phosphatidylcholine and transfer to phosphatidyl ethanolamine in homogenates of murine thymocytes, *J. Trotter, E. Ferber* 237
- A new type of ATP-activated bioluminescent system in the millipede *Lumino-desmus sequoiae*, *O. Shimomura* 242
- The photochemical quantum yield of bacteriorhodopsin is pH independent: A photoacoustic study, *M. Renard, M. Delmelle* 245
- Effect of Triton X-100 and of deuteration on the amplitude of the O<sub>640</sub>-intermediate in the bacteriorhodopsin photocycle, *W. Mantele, F. Siebert, W. Kreutz* 249
- Lysine 216 is a binding site of the retinyl moiety in bacteriorhodopsin, *H. D. Lemke, D. Oesterhelt* 255
- The isolated F<sub>0</sub> of *Escherichia coli* ATP-synthase is reconstitutively active in H<sup>+</sup>-conduction and ATP-dependent energy transduction, *P. Friedl, H. U. Schairer* 261
- Electric field-induced pK-changes in bacteriorhodopsin, *K. Tsuji, E. Neumann* 265
- Evidence for a calcium-gated cation channel in sarcoplasmic reticulum vesicles, *N. Bennett, Y. Dupont* 269
- Phosphate-independent calcium efflux from liver mitochondria, *F. Zoccarato, D. G. Nicholls* 275
- Restricted lateral diffusion of surface membrane components in *Tetrahymena thermophila*, *R. J. Hill* 278

|   |     |
|---|-----|
| The reconstitution of human C1, the first complement component: Binding of C1r and C1s to C1q influences the C1q conformation, <i>M. D. Golan, T. Hitschold, M. Loos</i>  | 281 |
| Binding of 5-chloropyrimidin-2-one to horse liver alcohol dehydrogenase, <i>J. Kovar, L. Skursky, S. G. Laland, R. Oftebro</i>  | 286 |
| Is the receptor-mediated endocytosis of cholera toxin a pre-requisite for its activation of adenylate cyclase in intact rat hepatocytes?, <i>M. D. Houslay, K. R. F. Elliott</i>  | 289 |
| Inactivation of $\gamma$ -cystathionase by $\gamma$ -fluorinated amino acids, <i>Th. A. Alston, H. Muramatsu, I. Ueda, H. J. Bright</i>   | 293 |
| 3'-Terminal nucleotide sequence of <i>Crithidia fasciculata</i> small ribosomal subunit RNA, <i>M. N. Schnare, M. W. Gray</i>   | 298 |
| Secreted proteins in the medium of microinjected <i>Xenopus</i> oocytes are degraded by oocyte proteases, <i>H. Soreq, R. Miskin</i>  | 305 |
| A heat-stable inhibitor protein for bovine brain cyclic nucleotide phosphodiesterase from <i>Escherichia coli</i> , <i>Y. Iwasa, K. Yonemitsu, K. Matsui, K. Fukunaga, E. Miyamoto</i>                                    | 311 |
| H1 and H5 histone arrangement in chromatin of pigeon erythrocytes, <i>V. A. Pospelov, A. M. Jerkin, A. T. Khachatryan</i>   | 315 |
| Anti-D-coated Rh-positive red cells will bind the first component of the complement pathway, C1q, <i>N. Hughes-Jones, S. Ghosh</i>  | 318 |
| Insulin stimulation of amino acid transport in primary cultured rat hepatocytes varies in direct proportion to insulin binding, <i>J. Dolais-Kitabgi, F. Checler, J.-F. Rey, O. Morin, P. Freychet</i>                    | 321 |
| A comparative study of mammalian neurophysin protein sequences, <i>D. H. Schlesinger, T. F. Audhya</i>  | 325 |
| Formation of novel hydroxylated eicosatetraenoic acids in preparations of human polymorphonuclear leukocytes, <i>J. A. Lindgren, G. Hansson, B. Samuelsson</i>  | 329 |
| Isoenzymes of rabbit slow myosin, <i>K. Inter, K. Mabuchi, F. A. Sreter</i>   | 336 |
| Production of hydroxyl radicals from paraquat radicals and $H_2O_2$ , <i>Ch. C. Winterbourn</i>   | 339 |
| Thiobarbituric acid-reactivity following iron-dependent free-radical damage to amino acids and carbohydrates, <i>J. M. C. Gutteridge</i>  | 343 |
| Formation of a thiobarbituric acid-reactive substance from deoxyribose in the presence of iron salts. The role of superoxide and hydroxyl radicals, <i>B. Halliwell, J. M. C. Gutteridge</i>                              | 347 |
| Yeast phosphoglycerate kinase purified by affinity elution has tightly bound 3-phosphoglycerate, <i>S. C. Condroy, B. Adams, R. H. Pain, L. A. Fothergill</i>   | 353 |
| The glutamate dehydrogenase system and the redox state of mitochondrial free nicotinamide adenine dinucleotide in myocardium, <i>E. M. Nuutinen, J. K. Hiltunen, I. E. Hassinen</i>                                       | 356 |
| The mechanism in the increase in renal ammoniogenesis in the rat with acute metabolic acidosis, <i>J. A. Lupianez, P. Hortelano, F. S. Medina, A. Sanchez-Pogo, N. McFarlane-Anderson, J. Barnswell, G. A. O. Alleyne</i> | 361 |
| Demonstration and partial characterization of endo-N-acetyl- $\beta$ -D-glucosaminidase in human tissues, <i>B. Overdijk, W. M. J. van der Kroef, J. J. W. Lisman, R. J. Pierce, J. Montreuil, G. Spik</i>                | 364 |
| Author Index  | 367 |
| Notes to Authors  | 372 |
| Table of Contents   | 373 |